

Colorful well-foundedness

Yann Pequignot

University of California, Los Angeles

YST workshop, Bernoulli Center, Lausanne
June 29, 2018

Part I
**Well-quasi-orders
and
Better-quasi-orders**

Well-quasi-orders

A **quasi-order** (qo) is a set Q together with a *reflexive* and *transitive* binary relation \leq .

Definition

A **well-quasi-order** (wqo) is a qo that satisfies one of the following equivalent conditions.

- 1 Q is well-founded and has no infinite antichain;

Well-quasi-orders

A **quasi-order** (qo) is a set Q together with a *reflexive* and *transitive* binary relation \leq .

Definition

A **well-quasi-order** (wqo) is a qo that satisfies one of the following equivalent conditions.

- 1 Q is well-founded and has no infinite antichain;
- 2 there exists no bad sequence, i.e. no $(q_n)_n$ s.t.

$$\forall m, n \in \omega \quad m < n \rightarrow q_m \not\leq q_n.$$

Well-quasi-orders

A **quasi-order** (qo) is a set Q together with a *reflexive* and *transitive* binary relation \leq .

Definition

A **well-quasi-order** (wqo) is a qo that satisfies one of the following equivalent conditions.

- 1 Q is well-founded and has no infinite antichain;
- 2 there exists no bad sequence, i.e. no $(q_n)_n$ s.t.

$$\forall m, n \in \omega \quad m < n \rightarrow q_m \not\leq q_n.$$

- 3 $\mathcal{P}(Q)$ is well-founded, under:

$$X \leq Y \iff \forall x \in X \exists y \in Y \ x \leq y.$$

Well-quasi-orders

Examples of wqos

- Finite quasi-orders

Well-quasi-orders

Examples of wqos

- Finite quasi-orders
- Well-orders

Well-quasi-orders

Examples of wqos

- Finite quasi-orders
- Well-orders
- If P and Q are wqo, then $P \times Q$ is wqo.

Well-quasi-orders

Examples of wqos

- Finite quasi-orders
- Well-orders
- If P and Q are wqo, then $P \times Q$ is wqo.
- (Higman 52') If P is wqo, then $P^{<\omega}$ is wqo under

$$(p_i)_{i < n} \leq (q_j)_{j < m} \iff \exists f : n \rightarrow m \text{ strictly increasing} \\ \text{s.t. } p_i \leq q_{f(i)} \text{ for all } i < n$$

Well-quasi-orders

Examples of wqos

- Finite quasi-orders
- Well-orders
- If P and Q are wqo, then $P \times Q$ is wqo.
- (Higman 52') If P is wqo, then $P^{<\omega}$ is wqo under

$$(p_i)_{i < n} \leq (q_j)_{j < m} \iff \exists f : n \rightarrow m \text{ strictly increasing} \\ \text{s.t. } p_i \leq q_{f(i)} \text{ for all } i < n$$

- (Laver 71') Countable linear orders under embeddability.

Well-quasi-orders


Examples of wqos

- Finite quasi-orders
- Well-orders
- If P and Q are wqo, then $P \times Q$ is wqo.
- (Higman 52') If P is wqo, then $P^{<\omega}$ is wqo under

$$(p_i)_{i < n} \leq (q_j)_{j < m} \iff \exists f : n \rightarrow m \text{ strictly increasing} \\ \text{s.t. } p_i \leq q_{f(i)} \text{ for all } i < n$$

- (Laver 71') Countable linear orders under embeddability.
- (Robertson-Seymour, 500 pages, 1983-2004) The finite undirected graphs under the minor relation.

A wqo Q such that $\mathcal{P}(Q)$ is not wqo



Rado's poset \mathcal{R}



Richard Rado, 1954.


\mathcal{R} is defined on $[\omega]^2$ by:

$$\{m_0, m_1\} \leq \{n_0, n_1\}$$

iff

$$\begin{cases} m_0 = n_0 \text{ and } m_1 \leq n_1, \text{ or} \\ m_0 < m_1 < n_0 < n_1 \end{cases}$$

A wqo Q such that $\mathcal{P}(Q)$ is not wqo



Rado's poset \mathcal{R}



Richard Rado, 1954.


\mathcal{R} is defined on $[\omega]^2$ by:

$$\{m_0, m_1\} \leq \{n_0, n_1\}$$

iff

$$\left\{ \begin{array}{l} m_0 = n_0 \text{ and } m_1 \leq n_1, \text{ or} \\ m_0 < m_1 < n_0 < n_1 \end{array} \right.$$

A wqo Q such that $\mathcal{P}(Q)$ is not wqo



Rado's poset \mathcal{R}



Richard Rado, 1954.


\mathcal{R} is defined on $[\omega]^2$ by:

$$\{m_0, m_1\} \leq \{n_0, n_1\}$$

iff

$$\begin{cases} m_0 = n_0 \text{ and } m_1 \leq n_1, \text{ or} \\ m_0 < m_1 < n_0 < n_1 \end{cases}$$

A wqo Q such that $\mathcal{P}(Q)$ is not wqo



Rado's poset \mathcal{R}



Richard Rado, 1954.


\mathcal{R} is defined on $[\omega]^2$ by:

$$\{m_0, m_1\} \leq \{n_0, n_1\}$$

iff

$$\left\{ \begin{array}{l} m_0 = n_0 \text{ and } m_1 \leq n_1, \text{ or} \\ m_0 < m_1 < n_0 < n_1 \end{array} \right.$$

A wqo Q such that $\mathcal{P}(Q)$ is not wqo



Richard Rado, 1954.

\mathcal{R} is defined on $[\omega]^2$ by:

$$\{m_0, m_1\} \leq \{n_0, n_1\}$$

iff

$$\left\{ \begin{array}{l} m_0 = n_0 \text{ and } m_1 \leq n_1, \text{ or} \\ m_0 < m_1 < n_0 < n_1 \end{array} \right.$$

Better quasi-orders

Fix a quasi-order Q and treat the element of Q as *atoms*, namely they have no elements but they are different from the empty set. We define by transfinite recursion:

$$\begin{aligned}Q_0^* &= Q \\ Q_{\alpha+1}^* &= \mathcal{P}^*(Q_\alpha^*) \quad (\text{the non-empty subsets of } V_\alpha^*) \\ Q_\lambda^* &= \bigcup_{\alpha < \lambda} Q_\alpha^*, \quad \text{for } \lambda \text{ limit.}\end{aligned}$$

Let

$$Q^* = \bigcup_{\alpha} Q_\alpha^*.$$

We define a quasi-order on Q^* via the existence of a winning strategy in a suitable game $G(X, Y)$.

Definition (Intuitive definition)


A quasi-order Q is a **better-quasi-order** if Q^* is well-founded.

Making sense of the definition

Let $(X_n)_n$ be a sequence in Q^* such that $m < n$ implies $X_n \not\leq X_m$.


Making sense of the definition

Let $(X_n)_n$ be a sequence in Q^* such that $m < n$ implies $X_n \not\leq X_m$.
For every infinite subset $N = \{n_0, n_1, n_2, \dots\}$ of ω , contemplate:



Making sense of the definition


Let $(X_n)_n$ be a sequence in Q^* such that $m < n$ implies $X_n \not\leq X_m$.
 For every infinite subset $N = \{n_0, n_1, n_2, \dots\}$ of ω , contemplate:



Where $S(N) = N \setminus \{\min N\}$. This defines a map $f : [\omega]^\infty \rightarrow Q$,
 $f(N) = Q_N$, such that $f(N) \not\leq f(S(N))$.

Making sense of the definition


Let $(X_n)_n$ be a sequence in Q^* such that $m < n$ implies $X_n \not\leq X_m$.
For every infinite subset $N = \{n_0, n_1, n_2, \dots\}$ of ω , contemplate:



Where $S(N) = N \setminus \{\min N\}$. This defines a map $f : [\omega]^\infty \rightarrow Q$,
 $f(N) = Q_N$, such that $f(N) \not\leq f(S(N))$.

Making sense of the definition


Let $(X_n)_n$ be a sequence in Q^* such that $m < n$ implies $X_n \not\leq X_m$.
 For every infinite subset $N = \{n_0, n_1, n_2, \dots\}$ of ω , contemplate:



Where $S(N) = N \setminus \{\min N\}$. This defines a map $f : [\omega]^\infty \rightarrow Q$,
 $f(N) = Q_N$, such that $f(N) \not\leq f(S(N))$.

Making sense of the definition


Let $(X_n)_n$ be a sequence in Q^* such that $m < n$ implies $X_n \not\leq X_m$.
 For every infinite subset $N = \{n_0, n_1, n_2, \dots\}$ of ω , contemplate:



Where $S(N) = N \setminus \{\min N\}$. This defines a map $f : [\omega]^\infty \rightarrow Q$,
 $f(N) = Q_N$, such that $f(N) \not\leq f(S(N))$.

Making sense of the definition


Let $(X_n)_n$ be a sequence in Q^* such that $m < n$ implies $X_n \not\leq X_m$.
 For every infinite subset $N = \{n_0, n_1, n_2, \dots\}$ of ω , contemplate:



Where $S(N) = N \setminus \{\min N\}$. This defines a map $f : [\omega]^\infty \rightarrow Q$,
 $f(N) = q_N$, such that $f(N) \not\leq f(S(N))$.

Making sense of the definition


Let $(X_n)_n$ be a sequence in Q^* such that $m < n$ implies $X_n \not\leq X_m$.
 For every infinite subset $N = \{n_0, n_1, n_2, \dots\}$ of ω , contemplate:



Where $S(N) = N \setminus \{\min N\}$. This defines a map $f : [\omega]^\infty \rightarrow Q$,
 $f(N) = q_N$, such that $f(N) \not\leq f(S(N))$.

Making sense of the definition


Let $(X_n)_n$ be a sequence in Q^* such that $m < n$ implies $X_n \not\leq X_m$.
 For every infinite subset $N = \{n_0, n_1, n_2, \dots\}$ of ω , contemplate:



Where $S(N) = N \setminus \{\min N\}$. This defines a map $f : [\omega]^\infty \rightarrow Q$,
 $f(N) = q_N$, such that $f(N) \not\leq f(S(N))$.

Making sense of the definition

Let $(X_n)_n$ be a sequence in Q^* such that $m < n$ implies $X_n \not\leq X_m$.
 For every infinite subset $N = \{n_0, n_1, n_2, \dots\}$ of ω , contemplate:




Where $S(N) = N \setminus \{\min N\}$. This defines a map $f : [\omega]^\infty \rightarrow Q$,
 $f(N) = q_N$, such that $f(N) \not\leq f(S(N))$.

Making sense of the definition

Let $(X_n)_n$ be a sequence in Q^* such that $m < n$ implies $X_n \not\leq X_m$.

For every infinite subset $N = \{n_0, n_1, n_2, \dots\}$ of ω , contemplate:




Where $S(N) = N \setminus \{\min N\}$. This defines a map $f : [\omega]^\infty \rightarrow Q$, $f(N) = Q_N$, such that $f(N) \not\leq f(S(N))$.

Making sense of the definition

Let $(X_n)_n$ be a sequence in Q^* such that $m < n$ implies $X_n \not\leq X_m$.

For every infinite subset $N = \{n_0, n_1, n_2, \dots\}$ of ω , contemplate:




Where $S(N) = N \setminus \{\min N\}$. This defines a map $f : [\omega]^\infty \rightarrow Q$, $f(N) = q_N$, such that $f(N) \not\leq f(S(N))$.

Making sense of the definition

Let $(X_n)_n$ be a sequence in Q^* such that $m < n$ implies $X_n \not\leq X_m$.

For every infinite subset $N = \{n_0, n_1, n_2, \dots\}$ of ω , contemplate:




Where $S(N) = N \setminus \{\min N\}$. This defines a map $f : [\omega]^\infty \rightarrow Q$, $f(N) = q_N$, such that $f(N) \not\leq f(S(N))$.

Making sense of the definition

Let $(X_n)_n$ be a sequence in Q^* such that $m < n$ implies $X_n \not\leq X_m$.

For every infinite subset $N = \{n_0, n_1, n_2, \dots\}$ of ω , contemplate:




Where $S(N) = N \setminus \{\min N\}$. This defines a map $f : [\omega]^\infty \rightarrow Q$, $f(N) = q_N$, such that $f(N) \not\leq f(S(N))$.

Making sense of the definition

Let $(X_n)_n$ be a sequence in Q^* such that $m < n$ implies $X_n \not\leq X_m$.


For every infinite subset $N = \{n_0, n_1, n_2, \dots\}$ of ω , contemplate:



Where $S(N) = N \setminus \{\min N\}$. This defines a map $f : [\omega]^\infty \rightarrow Q$, $f(N) = q_N$, such that $f(N) \not\leq f(S(N))$.

Making sense of the definition


Let $(X_n)_n$ be a sequence in Q^* such that $m < n$ implies $X_n \not\leq X_m$.
 For every infinite subset $N = \{n_0, n_1, n_2, \dots\}$ of ω , contemplate:



Where $S(N) = N \setminus \{\min N\}$. This defines a map $f : [\omega]^\infty \rightarrow Q$,
 $f(N) = Q_N$, such that $f(N) \not\leq f(S(N))$.

Making sense of the definition


Let $(X_n)_n$ be a sequence in Q^* such that $m < n$ implies $X_n \not\leq X_m$.
 For every infinite subset $N = \{n_0, n_1, n_2, \dots\}$ of ω , contemplate:



Where $S(N) = N \setminus \{\min N\}$. This defines a map $f : [\omega]^\infty \rightarrow Q$,
 $f(N) = q_N$, such that $f(N) \not\leq f(S(N))$.

Making sense of the definition


Let $(X_n)_n$ be a sequence in Q^* such that $m < n$ implies $X_n \not\leq X_m$.
 For every infinite subset $N = \{n_0, n_1, n_2, \dots\}$ of ω , contemplate:



Where $S(N) = N \setminus \{\min N\}$. This defines a map $f : [\omega]^\infty \rightarrow Q$,
 $f(N) = q_N$, such that $f(N) \not\leq f(S(N))$.

Making sense of the definition


Let $(X_n)_n$ be a sequence in Q^* such that $m < n$ implies $X_n \not\leq X_m$.
 For every infinite subset $N = \{n_0, n_1, n_2, \dots\}$ of ω , contemplate:



Where $S(N) = N \setminus \{\min N\}$. This defines a map $f : [\omega]^\infty \rightarrow Q$,
 $f(N) = q_N$, such that $f(N) \not\leq f(S(N))$.

Making sense of the definition


Let $(X_n)_n$ be a sequence in Q^* such that $m < n$ implies $X_n \not\leq X_m$.
 For every infinite subset $N = \{n_0, n_1, n_2, \dots\}$ of ω , contemplate:



Where $S(N) = N \setminus \{\min N\}$. This defines a map $f : [\omega]^\infty \rightarrow Q$,
 $f(N) = q_N$, such that $f(N) \not\leq f(S(N))$.

Making sense of the definition


Let $(X_n)_n$ be a sequence in Q^* such that $m < n$ implies $X_n \not\leq X_m$.
 For every infinite subset $N = \{n_0, n_1, n_2, \dots\}$ of ω , contemplate:



Where $S(N) = N \setminus \{\min N\}$. This defines a map $f : [\omega]^\infty \rightarrow Q$,
 $f(N) = q_N$, such that $f(N) \not\leq f(S(N))$.

Making sense of the definition


Let $(X_n)_n$ be a sequence in Q^* such that $m < n$ implies $X_n \not\leq X_m$.
 For every infinite subset $N = \{n_0, n_1, n_2, \dots\}$ of ω , contemplate:



Where $S(N) = N \setminus \{\min N\}$. This defines a map $f : [\omega]^\infty \rightarrow Q$,
 $f(N) = q_N$, such that $f(N) \not\leq f(S(N))$.

Making sense of the definition

Let $(X_n)_n$ be a sequence in Q^* such that $m < n$ implies $X_n \not\leq X_m$.
 For every infinite subset $N = \{n_0, n_1, n_2, \dots\}$ of ω , contemplate:



Where $S(N) = N \setminus \{\min N\}$. This defines a map $f : [\omega]^\infty \rightarrow Q$,
 $f(N) = q_N$, such that $f(N) \not\leq f(S(N))$.

A working definition for better-quasi-orders



$[\omega]^\infty$ is a Polish space homeomorphic to ω^ω .
Considering Q with the discrete topology, we just proved:

Proposition

If Q^* is *ill-founded*, then there exists a continuous map $f : [\omega]^\infty \rightarrow Q$ s.t. $f(N) \not\leq f(S(N))$ for every N .

And in fact, this is an equivalence. So we get:

Definition (Working definition, Nash-Williams 65')

A quasi-order Q is a *better-quasi-order* (bqo) if for every continuous map $f : [\omega]^\infty \rightarrow Q$ there exists $N \in [\omega]^\infty$ such that $f(N) \leq f(S(N))$.

Examples of better-quasi-orders

Theorem (Nash-Williams, Galvin-Prikry)

For every finite partition $[\omega]^\infty = B_0 \cup \dots \cup B_n$ into Borel sets, there exists an infinite $X \subseteq \omega$ such that $[X]^\infty \subseteq B_i$ for some $i \in \{0, \dots, n\}$.

Examples of bqos

- Finite quasi-orders

Examples of better-quasi-orders

Theorem (Nash-Williams, Galvin-Prikry)

For every finite partition $[\omega]^\infty = B_0 \cup \dots \cup B_n$ into Borel sets, there exists an infinite $X \subseteq \omega$ such that $[X]^\infty \subseteq B_i$ for some $i \in \{0, \dots, n\}$.

Examples of bqos

- Finite quasi-orders
- Well-orders

Examples of better-quasi-orders

Theorem (Nash-Williams, Galvin-Prikry)

For every finite partition $[\omega]^\infty = B_0 \cup \dots \cup B_n$ into Borel sets, there exists an infinite $X \subseteq \omega$ such that $[X]^\infty \subseteq B_i$ for some $i \in \{0, \dots, n\}$.

Examples of bqos

- Finite quasi-orders
- Well-orders
- If P and Q are bqo, then $P \times Q$ is bqo.

Examples of better-quasi-orders

Theorem (Nash-Williams, Galvin-Prikry)

For every finite partition $[\omega]^\infty = B_0 \cup \dots \cup B_n$ into Borel sets, there exists an infinite $X \subseteq \omega$ such that $[X]^\infty \subseteq B_i$ for some $i \in \{0, \dots, n\}$.

Examples of bqos

- Finite quasi-orders
- Well-orders
- If P and Q are bqo, then $P \times Q$ is bqo.
- (Nash-Williams) If P is bqo, then P^ω is bqo under

$$(p_i)_{i \in \omega} \leq (q_j)_{j \in \omega} \iff \exists f : \omega \rightarrow \omega \text{ strictly increasing} \\ \text{s.t. } p_i \leq q_{f(i)} \text{ for all } i \in \omega$$

Examples of better-quasi-orders

Theorem (Nash-Williams, Galvin-Prikry)

For every finite partition $[\omega]^\infty = B_0 \cup \dots \cup B_n$ into Borel sets, there exists an infinite $X \subseteq \omega$ such that $[X]^\infty \subseteq B_i$ for some $i \in \{0, \dots, n\}$.

Examples of bqos

- Finite quasi-orders
- Well-orders
- If P and Q are bqo, then $P \times Q$ is bqo.
- (Nash-Williams) If P is bqo, then P^ω is bqo under

$$(p_i)_{i \in \omega} \leq (q_j)_{j \in \omega} \iff \exists f : \omega \rightarrow \omega \text{ strictly increasing} \\ \text{s.t. } p_i \leq q_{f(i)} \text{ for all } i \in \omega$$

- (Laver 71') Countable linear orders under embeddability.

Examples of better-quasi-orders

Theorem (Nash-Williams, Galvin-Prikry)

For every finite partition $[\omega]^\infty = B_0 \cup \dots \cup B_n$ into Borel sets, there exists an infinite $X \subseteq \omega$ such that $[X]^\infty \subseteq B_i$ for some $i \in \{0, \dots, n\}$.

Examples of bqos

- Finite quasi-orders
- Well-orders
- If P and Q are bqo, then $P \times Q$ is bqo.
- (Nash-Williams) If P is bqo, then P^ω is bqo under

$$(p_i)_{i \in \omega} \leq (q_j)_{j \in \omega} \iff \exists f : \omega \rightarrow \omega \text{ strictly increasing} \\ \text{s.t. } p_i \leq q_{f(i)} \text{ for all } i \in \omega$$

- (Laver 71') Countable linear orders under embeddability.
- every "naturally occurring" wqo.

Part II
Infinite vs. infinite
Borel chromatic number

Directed graphs

- A *directed graph* is a pair $\mathbf{D} = (X, D)$ where D is an irreflexive binary relation on X .
- A *homomorphism* from (X, D) to (X', D') is a map $h : X \rightarrow X'$ such that $x D y$ implies $h(x) D' h(y)$.
- A *coloring* of a directed graph (X, D) is a map $c : X \rightarrow Y$ such that $x D x'$ implies $c(x) \neq c(x')$.
- The *chromatic number* of $\mathbf{D} = (X, D)$, $\chi(\mathbf{D})$, is the smallest cardinality of a set Y s.t. there exists a coloring $c : X \rightarrow Y$.

Borel chromatic number

- If X is a Polish space, the *Borel chromatic number*, $\chi_B(\mathbf{D})$, of $\mathbf{D} = (X, D)$ is the smallest cardinality of a Polish space Y such that there exists a Borel coloring $c : X \rightarrow Y$.
- Write $(X, D) \preceq (X', D')$, (\preceq_C, \preceq_B) if there exists a (continuous, Borel) homomorphism from (X, D) to (X', D') .

Remark:

- 1 $\chi_B(\mathbf{D}) \in \{1, 2, 3, \dots, \aleph_0, 2^{\aleph_0}\}$.
- 2 if $(X, D) \preceq_B (X', D')$ then $\chi_B(X, D) \leq \chi_B(X', D')$.

Borel chromatic number

- If X is a Polish space, the *Borel chromatic number*, $\chi_B(\mathbf{D})$, of $\mathbf{D} = (X, D)$ is the smallest cardinality of a Polish space Y such that there exists a Borel coloring $c : X \rightarrow Y$.
- Write $(X, D) \preceq (X', D')$, (\preceq_c, \preceq_B) if there exists a (continuous, Borel) homomorphism from (X, D) to (X', D') .

1 $\chi_B(\mathbf{D}) \in \{1, 2, 3, \dots, \aleph_0, 2^{\aleph_0}\}$.

Remark:

2 if $(X, D) \preceq_B (X', D')$ then $\chi_B(X, D) \leq \chi_B(X', D')$.

Theorem (Kechris–Solecki–Todorćević, 96)

There is a graph \mathbf{G}_0 on 2^ω s.t. for every analytic graph $\mathbf{G} = (X, G)$ on a Polish space X , exactly one of the following holds:

1 $\chi_B(\mathbf{G}) \leq \aleph_0$,

2 $\mathbf{G}_0 \preceq_c \mathbf{G}$ (and therefore $\chi_B(\mathbf{G}) = 2^{\aleph_0}$).

Graphs generated by a function

For any function $f : X \rightarrow X$, let (X, f) denote the directed graph whose arrows are given by:

$$x D_f y \leftrightarrow x \neq y \text{ and } f(x) = y.$$

Graphs generated by a function

For any function $f : X \rightarrow X$, let (X, f) denote the directed graph whose arrows are given by:

$$x D_f y \leftrightarrow x \neq y \text{ and } f(x) = y.$$

Remark: If X is Polish and f is Borel, then $\chi_B(X, f) \leq \aleph_0$.

Theorem (Kechris–Solecki–Todorćević,)

Let $f : X \rightarrow X$ be a Borel function with no fixed point. Then the following are equivalent:

- 1** $\chi_B(X, f) \leq 3$,
- 2** $\chi_B(X, f)$ is finite,

Graphs generated by a function

For any function $f : X \rightarrow X$, let (X, f) denote the directed graph whose arrows are given by:

$$x D_f y \leftrightarrow x \neq y \text{ and } f(x) = y.$$

Remark: If X is Polish and f is Borel, then $\chi_B(X, f) \leq \aleph_0$.

Theorem (Kechris–Solecki–Todorćević, Miller)

Let $f : X \rightarrow X$ be a Borel function with no fixed point. Then the following are equivalent:

- 1** $\chi_B(X, f) \leq 3$,
- 2** $\chi_B(X, f)$ is finite,
- 3** there exists a Borel subset B of X such that

$$\forall x \in X \ (\exists^\infty m \in \omega \ f^m(x) \in B \text{ and } \exists^\infty n \in \omega \ f^n(x) \notin B).$$

Finite vs Infinite: The shift graph (again :-)

Let $[\omega]^\infty$ be the space of infinite subsets of ω . As a subspace of 2^ω it is Polish and homeomorphic to ω^ω . The *shift map* is defined by

$$\begin{aligned} S : [\omega]^\infty &\longmapsto [\omega]^\infty \\ X &\longrightarrow X \setminus \{\min X\} \end{aligned}$$

The *Shift Graph* is the directed graph $\mathcal{G}_S = ([\omega]^\infty, S)$.

Finite vs Infinite: The shift graph (again :-)

Let $[\omega]^\infty$ be the space of infinite subsets of ω . As a subspace of 2^ω it is Polish and homeomorphic to ω^ω . The *shift map* is defined by

$$\begin{aligned} S : [\omega]^\infty &\longmapsto [\omega]^\infty \\ X &\longrightarrow X \setminus \{\min X\} \end{aligned}$$

The *Shift Graph* is the directed graph $\mathcal{G}_S = ([\omega]^\infty, S)$.

- As \mathcal{G}_S is acyclic, we have $\chi(\mathcal{G}_S) = 2$ (Axiom of choice :-)

Finite vs Infinite: The shift graph (again :-)

Let $[\omega]^\infty$ be the space of infinite subsets of ω . As a subspace of 2^ω it is Polish and homeomorphic to ω^ω . The *shift map* is defined by

$$\begin{aligned} S : [\omega]^\infty &\longmapsto [\omega]^\infty \\ X &\longrightarrow X \setminus \{\min X\} \end{aligned}$$

The *Shift Graph* is the directed graph $\mathcal{G}_S = ([\omega]^\infty, S)$.

- As \mathcal{G}_S is acyclic, we have $\chi(\mathcal{G}_S) = 2$ (Axiom of choice :-)
- The Galvin–Prikry theorem: for every finite Borel coloring of $[\omega]^\infty$ there exists an infinite $X \subseteq \omega$ such that $[X]^\infty$ is monochromatic. In particular X and $S(X)$ have same color. Hence $\chi_B(\mathcal{G}_S)$ is infinite. But $c : [\omega]^\infty \rightarrow \omega$, $X \mapsto \min X$ is a continuous coloring, so

$$\chi_B(\mathcal{G}_S) = \aleph_0.$$

Finite vs infinite

(Kechris–Solecki–Todorćević, 1996) Is the following true?

If X is a Polish space and $f : X \rightarrow X$ is a Borel function, then exactly one of the following holds:

- 1 The Borel chromatic number of (X, f) is finite;
- 2 $\mathcal{G}_S \preceq_c (X, f)$.

Finite vs infinite

(Kechris–Solecki–Todorćević, 1996) Is the following true?

If X is a Polish space and $f : X \rightarrow X$ is a Borel function, then exactly one of the following holds:

- 1 The Borel chromatic number of (X, f) is finite;
- 2 $\mathcal{G}_S \preceq_c (X, f)$.

The answer is negative.

Finite vs infinite

Theorem (P)

There exists a closed subset C of $[\omega]^\infty$ such that

- *$X \in C$ implies $S(X) \in C$,*
- *the Borel chromatic number of (C, S) is infinite,*
- *there is no Borel homomorphism from \mathcal{G}_S to (C, S) .*

- However no “natural” example is known.
- The proof consists of showing that the collection of closed sets as above is a true Π_2^1 set, hence non empty.
- It relies on a representation theorem for Σ_2^1 sets.

Finite vs infinite

Theorem (P)

There exists a closed subset C of $[\omega]^\infty$ such that

- *$X \in C$ implies $S(X) \in C$,*
- *the Borel chromatic number of (C, S) is infinite,*
- *there is no Borel homomorphism from \mathcal{G}_S to (C, S) .*

- However no “natural” example is known.
- The proof consists of showing that the collection of closed sets as above is a true Π_2^1 set, hence non empty.
- It relies on a representation theorem for Σ_2^1 sets.

Actually there is no basis result at all since:

Theorem (Todorčević, Vidnyánszky)

The set of codes for closed subsets C of $[\omega]^\infty$ for which (C, S) has finite Borel chromatic number is Σ_2^1 -complete.

Representation of analytic sets

Let $\mathbb{G} = 2^\omega$ be the Polish space of (codes for) countable directed graphs, where $\alpha \in 2^\omega$ codes (X_α, D_α) given by

$$X_\alpha = \{n \mid \alpha(\langle n, n \rangle) = 0\}, \text{ and}$$

$$m D_\alpha n \leftrightarrow \alpha(\langle m, n \rangle) = 1 \text{ and } m, n \in X_\alpha.$$

Representation of analytic sets

Let $\mathbb{G} = 2^\omega$ be the Polish space of (codes for) countable directed graphs, where $\alpha \in 2^\omega$ codes (X_α, D_α) given by

$$X_\alpha = \{n \mid \alpha(\langle n, n \rangle) = 0\}, \text{ and} \\ m D_\alpha n \leftrightarrow \alpha(\langle m, n \rangle) = 1 \text{ and } m, n \in X_\alpha.$$

Proposition (Folklore)

A subset A of ω^ω is Σ_1^1 iff there exists a continuous function $\omega^\omega \rightarrow \mathbb{G}$, $\alpha \mapsto \mathbf{G}_\alpha$ such that

$$\alpha \in A \iff (\omega, <) \preceq \mathbf{G}_\alpha$$

Proof sketch: Let $T = \{(x \upharpoonright_n, y \upharpoonright_n) \mid (x, y) \in C \text{ and } n \in \omega\}$ and set $T(\alpha) = \{s \in \omega^{<\omega} \mid (\alpha \upharpoonright_s, s) \in T\}$. We have

$$\begin{aligned} \alpha \in A &\iff \exists \beta \in \omega^\omega \forall n (\alpha \upharpoonright_n, \beta \upharpoonright_n) \in T \\ &\iff \exists \beta \in \omega^\omega \forall n \beta \upharpoonright_n \in T(\alpha) \\ &\iff (\omega, <) \preceq (T(\alpha), \sqsubset) = \mathbf{G}_\alpha. \end{aligned}$$

Representation of Σ_2^1 sets

Recall that a subset $P \subseteq \omega^\omega$ is Σ_2^1 if there exists a closed subset C of $\omega^\omega \times \omega^\omega \times \omega^\omega$ such that

$$\alpha \in P \iff \exists \beta \in \omega^\omega \forall \gamma \in \omega^\omega (\alpha, \beta, \gamma) \notin C.$$

Representation of Σ_2^1 sets

Recall that a subset $P \subseteq \omega^\omega$ is Σ_2^1 if there exists a closed subset C of $\omega^\omega \times \omega^\omega \times \omega^\omega$ such that

$$\alpha \in P \iff \exists \beta \in \omega^\omega \forall \gamma \in \omega^\omega (\alpha, \beta, \gamma) \notin C.$$

Theorem (Marcone, 95')

A subset $P \subseteq \omega^\omega$ is Σ_2^1 iff there exists a continuous function $\omega^\omega \rightarrow \mathbb{G}$, $\alpha \mapsto \mathbf{G}_\alpha$ such that

$$\alpha \in P \iff \mathcal{G}_S \preceq_c \mathbf{G}_\alpha.$$

Again any $\mathbf{G} \in \mathbb{G}$ is considered with the discrete topology.

A Π_2^1 complete set

Corollary

$\{\mathbf{G} \in \mathbb{G} \mid \mathcal{G}_S \preceq_c \mathbf{G}\}$ is a Σ_2^1 non Π_2^1 subset of \mathbb{G} .

Proof.

- It is not too hard to give a Σ_2^1 definition.
- Suppose it is also Π_2^1 . As Π_2^1 is closed under continuous preimages, the representation theorem implies that $\Sigma_2^1 \subseteq \Pi_2^1$. This would contradict the existence of a universal Σ_2^1 set. \square

Definition

A countable directed graph $\mathbf{G} \in \mathbb{G}$ is **better** if $\mathcal{G}_S \not\preceq_c \mathbf{G}$ when the vertex set is considered with the discrete topology.

The set

$$BG = \{\mathbf{G} \in \mathbb{G} \mid \mathcal{G}_S \not\preceq_c \mathbf{G}\}$$

of better graphs is a Π_2^1 -complete set. In particular, not Σ_2^1 .

Shift on rays of a countable directed graph

Let $\mathbf{G} = (X, D)$ be directed graph on $X \subseteq \omega$.

Define the *Ray Graph* of \mathbf{G} as the directed graph $(\vec{\mathbf{G}}, S)$ where:

$$\vec{\mathbf{G}} = \{(n_i)_{i \in \omega} \in X^\omega \mid \forall i \in \omega \ n_i D n_{i+1}\}$$

and the shift map $S : \vec{\mathbf{G}} \rightarrow \vec{\mathbf{G}}$ given by

$$S((n_i)_{i \in \omega}) = (n_{i+1})_{i \in \omega}.$$

- If $\mathbf{G} = (\omega, <)$, then $\vec{\mathbf{G}} = [\omega]^\infty$.
- If $\mathbf{G} = (\omega, s)$, $s : n \mapsto n + 1$, then $\vec{\mathbf{G}} = \{\omega \setminus n \mid n \in \omega\}$.

Shift on rays of a countable directed graph

Let $\mathbf{G} = (X, D)$ be directed graph on $X \subseteq \omega$.

Define the *Ray Graph* of \mathbf{G} as the directed graph $(\vec{\mathbf{G}}, S)$ where:

$$\vec{\mathbf{G}} = \{(n_i)_{i \in \omega} \in X^\omega \mid \forall i \in \omega \ n_i D n_{i+1}\}$$

and the shift map $S : \vec{\mathbf{G}} \rightarrow \vec{\mathbf{G}}$ given by

$$S((n_i)_{i \in \omega}) = (n_{i+1})_{i \in \omega}.$$

- If $\mathbf{G} = (\omega, <)$, then $\vec{\mathbf{G}} = [\omega]^\infty$.
- If $\mathbf{G} = (\omega, s)$, $s : n \mapsto n + 1$, then $\vec{\mathbf{G}} = \{\omega \setminus n \mid n \in \omega\}$.

Proposition

For every $\mathbf{G} \in \mathbb{G}$:

$$\begin{aligned} \mathcal{G}_S \preceq_c \mathbf{G} &\iff \mathcal{G}_S \preceq_c (\vec{\mathbf{G}}, S), \\ &\iff \mathcal{G}_S \preceq_B (\vec{\mathbf{G}}, S). \end{aligned}$$

A very discrete graph

Recall that $BG = \{\mathbf{G} \in \mathbb{G} \mid \mathcal{G}_S \not\leq_c \mathbf{G}\}$ is Π_2^1 -complete.

Theorem

There exists $\mathbf{G} \in \mathbb{G}$ such that

$$\chi_B(\vec{\mathbf{G}}, S) = \aleph_0 \quad \text{and} \quad \mathcal{G}_S \not\leq_B (\vec{\mathbf{G}}, S).$$

Sketch of the proof.

- Prove that the set $\tilde{F} = \{\mathbf{G} \in \mathbb{G} \mid \chi_B(\vec{\mathbf{G}}) < \aleph_0\}$ is Σ_2^1 .
- Notice that $\tilde{F} \subseteq BG$: for if $\mathcal{G}_S \leq_c \mathbf{G}$, then $\mathcal{G}_S \leq_c \vec{\mathbf{G}}$ and so $\aleph_0 = \chi_B(\mathcal{G}_S) \leq \chi_B(\vec{\mathbf{G}})$.
- Since BG is not Σ_2^1 , we cannot have $\tilde{F} = BG$. Hence there exists \mathbf{G} with $\mathbf{G} \in BG$ and $\mathbf{G} \notin \tilde{F}$. Such a \mathbf{G} is as desired. \square

Part III
Ordering functions

Ordering functions

- One way to understand objects consists of ordering them.
- For sets $A, B \subseteq \omega^\omega$, continuous reducibility (Wadge \leq):

$$A \leq_W B \iff \exists f : \omega^\omega \rightarrow \omega^\omega \text{ continuous such that} \\ \forall x \in \omega^\omega (x \in A \leftrightarrow f(x) \in B).$$

- For equivalence relations E, F on ω^ω , Borel reducibility:

$$E \leq_B F \iff \exists f : \omega^\omega \rightarrow \omega^\omega \text{ Borel such that} \\ \forall x, y \in \omega^\omega (x E y \leftrightarrow f(x) F f(y)).$$

- What about functions?

Ordering functions

- One way to understand objects consists of ordering them.
- For sets $A, B \subseteq \omega^\omega$, continuous reducibility (Wadge \leq):

$$A \leq_W B \iff \exists f : \omega^\omega \rightarrow \omega^\omega \text{ continuous such that} \\ \forall x \in \omega^\omega (x \in A \leftrightarrow f(x) \in B).$$

- For equivalence relations E, F on ω^ω , Borel reducibility:

$$E \leq_B F \iff \exists f : \omega^\omega \rightarrow \omega^\omega \text{ Borel such that} \\ \forall x, y \in \omega^\omega (x E y \leftrightarrow f(x) F f(y)).$$

- What about functions?

All spaces considered are Polish zero-dimensional spaces, denoted by variables X, Y, \dots

Continuous reducibility on functions

Definition (Hertling-Weihrach, Carroy)

Say that $f : X \rightarrow Y$ reduces to $g : X' \rightarrow Y'$ if there are $\sigma : X \rightarrow X'$ continuous and $\tau : \text{im}(g \circ \sigma) \rightarrow Y$ continuous such that $f = \tau \circ g \circ \sigma$.

$$\begin{array}{ccc} X' & \xrightarrow{g} & Y' \\ \sigma \uparrow & \Downarrow & \downarrow \tau \\ X & \xrightarrow{f} & Y \end{array}$$

Theorem (Carroy, 2012)

Continuous reducibility is a well-order on *continuous* functions with *compact* domains.


Conjecture (Carroy)

Continuous reducibility is a wqo on *continuous* functions.

Topological embeddability on functions

Definition

Say that $f : X \rightarrow Y$ embeds into $g : X' \rightarrow Y'$ if there are embeddings $\sigma : X \rightarrow X'$ and $\tau : \text{im } f \rightarrow Y'$ such that $\tau \circ f = g \circ \sigma$.



- Embeddability is finer than reducibility: $f \sqsubseteq g \rightarrow f \leq g$.
- The projection $p : \omega^\omega \times \omega^\omega \rightarrow \omega^\omega$ is a **maximum** for continuous functions: $f : X \rightarrow Y$ is continuous iff $f \sqsubseteq p$.
- The two discontinuous functions

$$d_0 : \omega + 1 \longrightarrow 2$$

$$\omega \longmapsto 0$$

$$n \longmapsto 1$$

$$d_1 : \omega + 1 \longrightarrow \omega$$

$$\omega \longmapsto 0$$

$$n \longmapsto n + 1$$

form a 2-element **basis** for discontinuous functions:
 $f : X \rightarrow Y$ is discontinuous iff $d_0 \sqsubseteq f$ or $d_1 \sqsubseteq f$.

Topological embeddability on functions, continued.

Theorem

The following classes admits a **minimum** under embeddability:

- 1 (Solecki, 98') The class of Baire class 1 functions that are not σ -continuous.
- 2 (Zapletal, 04') The Borel functions that are not σ -continuous.
- 3 (Carroy-Miller, 17') The class of Baire class 1 functions that are not F_σ -to-one.

Theorem (Carroy-Miller, 17')

The following classes admits a **finite basis** under embeddability:

- 1 The Borel functions that are not in the first Baire class.
- 2 The Borel functions that are not σ -continuous with closed witnesses.

Conjecture, $\alpha > 1$:

The Borel functions that are not Baire class α admit a finite basis.

Order and Chaos

For X compact, $C(X, Y)$ denotes the space of continuous functions $X \rightarrow Y$ with the topology of uniform convergence.

Proposition (Carroy, P., Vidnyánszky)

If X, Y are Polish and X is compact, then embeddability is an analytic quasi-order on $C(X, Y)$.

An analytic qo Q on a Polish space Z is **analytic complete** if it Borel reduces every analytic qo on any Polish space.

Order and Chaos

For X compact, $C(X, Y)$ denotes the space of continuous functions $X \rightarrow Y$ with the topology of uniform convergence.

Proposition (Carroy, P., Vidnyánszky)

If X, Y are Polish and X is compact, then embeddability is an analytic quasi-order on $C(X, Y)$.

An analytic qo Q on a Polish space Z is **analytic complete** if it Borel reduces every analytic qo on any Polish space.

Theorem (Carroy, P., Vidnyánszky)

Suppose that X, Y are Polish zero-dimensional and X is compact. Then exactly one of the following holds:

- 1** *embeddability on $C(X, Y)$ is an analytic complete quasi-order,*
- 2** *embeddability on $C(X, Y)$ is a wqo. In fact, a bqo.*

*Moreover **1** holds exactly when X has infinitely many non-isolated points and Y is not discrete. For instance for $C(2^\omega, 2^\omega)$.*

Chaos

Let \mathbb{G} denote the Polish space of (simple) undirected graphs with vertex set \mathbb{N} .

For $G, H \in \mathbb{G}$ let

$G \leq_i H \iff$ there is an injective homomorphism from G to H .

Theorem (Louveau-Rosendal)

The $qo \leq_i$ on \mathbb{G} is an analytic complete quasi-order.

Theorem (Carroy, P., Vidnyánszky)

There is a continuous function $\mathbb{G} \rightarrow C(\omega^2 + 1, \omega + 1)$, $G \mapsto f^G$ that reduces \leq_i to \sqsubseteq :

$$G \leq_i H \iff f^G \sqsubseteq f^H.$$

So embeddability on $C(\omega^2 + 1, \omega + 1)$ is an analytic complete qo .

Order

Let \mathbb{Q} be the space of rationals, (P, \leq_P) a quasi-order.

Let $P^{\mathbb{Q}}$ be the set of maps $l : \mathbb{Q} \rightarrow P$ quasi-ordered by

$l_0 \leq l_1 \iff$ there is a topological embedding $\tau : \mathbb{Q} \rightarrow \mathbb{Q}$
such that $l_0(q) \leq_P l_1(\tau(q))$ for all $q \in \mathbb{Q}$.

Theorem (van Engelen-Miller-Steel)

If P is bqo, then $P^{\mathbb{Q}}$ is bqo.

Theorem (van Engelen-Miller-Steel, Carroy)

The Polish 0-dimensional spaces with embeddability are bqo.

Proposition (Carroy, P., Vidnyánszky)

The locally constant maps are bqo under embeddability.

In search of a specific example

Consider the set $2^{<\omega}$ of finite binary words equipped with the subword ordering, i.e.

$u \preceq v \iff$ there exists a strictly increasing map $h : |u| \rightarrow |v|$
such that for every $i < |u|$ we have $u(i) = v(h(i))$,

where $|u|$ denotes the length of $u \in 2^{<\omega}$. E.g. $01 \preceq 100100$.

- Let $\mathbf{H} = (2^{<\omega}, H)$ where $u H v \iff u \preceq v$.
- Since $(2^{<\omega}, \preceq)$ is a *better-quasi-order*, so $\mathcal{G}_S \not\prec_c (2^{<\omega}, H)$ and so $\mathcal{G}_S \not\prec_c \vec{H}$.

Question

What is the Borel chromatic number of \vec{H} ? Is it \aleph_0 ?

Remark: there is no **continuous** 2-coloring of \vec{H} .